

**COSEWIC**  
**Assessment and Status Report**

on the


**Golden-winged Warbler**  
*Vermivora chrysoptera*

in Canada



**THREATENED**  
**2006**

**COSEWIC**  
COMMITTEE ON THE STATUS OF  
ENDANGERED WILDLIFE  
IN CANADA



**COSEPAC**  
COMITÉ SUR LA SITUATION  
DES ESPÈCES EN PÉRIL  
AU CANADA



©





## COSEWIC Assessment Summary

### Assessment Summary – April 2006

**Common name**

Golden-winged Warbler

**Scientific name**

*Vermivora chrysoptera*

**Status**

Threatened

**Reason for designation**

This small songbird has declined by 79% over the last 10 years according to Breeding Bird Survey data from Canada. The main threat appears to be competition and genetic swamping (hybridization) from the closely related Blue-winged Warbler, which is spreading north because of habitat change and perhaps climate change.

**Occurrence**

Manitoba, Ontario, Quebec

**Status history**

Designated Threatened in April 2006. Assessment based on a new status report.



The species is known to be strictly insectivorous during the breeding season. The preferred diet consists mainly of tortricid moths and their larvae, although other moths and their pupae, other winged insects and spiders are sometimes consumed. Similar feeding habits are also seen on the wintering grounds.

### **Population sizes and trends**

The Golden-winged Warbler has been experiencing population declines for at least 30 years and is currently one of the fastest declining passerine species in North America. In Canada, the species was showing population growth

*Vermivora* genus (one of which, the Bachman's Warbler, is extinct). The Golden-winged Warbler can be seen as a representative for many other successional scrub breeding birds that are facing widespread population declines. As such, the conservation of this species will benefit numerous other avian species in Canada.

### **Existing protection**

The Golden-winged Warbler is not federally listed in the United States, although it has designation in 12 states and is currently under federal status assessment in the U.S. to determine whether it warrants protection under the U.S. Endangered Species Act. In Québec its status is "Likely to be designated as threatened or vulnerable". The Migratory Bird Treaty Act of 1918 provides protection from taking, killing or possessing the species. NatureServe gives the Golden-winged Warbler a global ranking of G4 (apparently secure – uncommon but not rare).

## **COSEWIC HISTORY**

The Committee on the Status of Endangered Wildlife in Canada (COSEWIC) was created in 1977 as a result of a recommendation at the Federal-Provincial Wildlife Conference held in 1976. It arose from the need for a single, official, scientifically sound, national listing of wildlife species at risk. In 1978, COSEWIC designated its first species and produced its first list of Canadian species at risk. Species designated at meetings of the full committee are added to the list. On June 5<sup>th</sup> 2003, the *Species at Risk Act* (SARA) was proclaimed. SARA establishes COSEWIC as an advisory body ensuring that species will continue to be assessed under a rigorous and independent scientific process.

## **COSEWIC MANDATE**

The Committee on the Status of Endangered Wildlife in Canada (COSEWIC) assesses the national status of wild species, subspecies, varieties, or other designatable units that are considered to be at risk in Canada. Designations are made on native species for the following taxonomic groups: mammals, birds, reptiles, amphibians, fishes, arthropods, molluscs, vascular plants, mosses, and lichens.

## **COSEWIC MEMBERSHIP**

COSEWIC comprises members from each provincial and territorial government wildlife agency, four federal entities (Canadian Wildlife Service, Parks Canada Agency, Department of Fisheries and Oceans, and the Federal Biodiversity Information Partnership, chaired by the Canadian Museum of Nature), three non-government science members and the co-chairs of the species specialist subcommittees and the Aboriginal Traditional Knowledge subcommittee. The Committee meets to consider status reports on candidate species.

## **DEFINITIONS (2006)**

# **COSEWIC Status Report**

on the

## **Golden-winged Warbler**

*Vermivora chrysoptera*

**in Canada**

2006



# TABLE OF CONTENTS

SPECIES INFORMATION.....	4
Name and classification.....	4
Morphological description.....	4
Genetic description.....	4
DISTRIBUTION.....	5
Global range.....	5
Canadian range.....	6
HABITAT.....	7
Habitat requirements.....	7
Habitat trends.....	8
Habitat protection/owners.....	8

S I O L O G Y

Table 2. State summaries of legal status of Golden-winged Warbler .....	19
Table 3. NatureServe ranks for the Golden-winged Warbler in the United States and Canada, by state/province .....	19

## SPECIES INFORMATION

### Name and classification

English name: Golden-winged Warbler  
Scientific name: *Vermivora chrysoptera* Linnaeus  
French name: Paruline à ailes dorées

### Morphological description

The Golden-winged Warbler is a small (9-11 g) wood warbler. Both sexes of adults have white underparts, grey back and wings, and a patch of yellow on each wing. Males have a striking black throat and eye patch which is a soft grey in females (Figure 1). Both sexes also have a yellow crown patch, although this tends to be much brighter in after second-year males than in second year males and females.



Figure 1. Male (left) and female (right) Golden-winged Warblers (photos by Rachel Fraser)

These birds could initially be confused with the Black-capped Chickadee (*Poecile atricapillus*; Confer 1992) that exhibits the same black throat patch and foraging style. However, the smaller size as well as the yellow patches on the wings and crown can easily distinguish them. If seen, it is unlikely that the Golden-winged Warbler would be confused with any other wood warbler species, except Golden-winged X Blue-winged Warbler hybrids (see Interspecific interactions section below). Identification by song can be more difficult, especially when considering hybrids.

### Genetic description

Golden-winged Warblers belong to the order Passeriformes (Perching Birds) and the Family Parulidae (Wood Warblers). Currently Golden-winged Warblers are considered members of the *Vermivora* genus that contains a total of nine species in North America. Recent phylogenetic work suggests that the genus *Vermivora* may be polyphyletic; Golden-winged Warblers and Blue-winged Warblers are not closely allied

to any other *Vermivora*, except perhaps the extinct Bachman's Warbler (*V. bachmanii*; I. Lovette, pers. comm.). These three species may be the only true *Vermivora* species, as the Blue-winged Warbler is the type-species for the genus (Klein *et al.* 2004).

## DISTRIBUTION

### Global range

Populations of breeding Golden-winged Warblers are found in northeastern regions of the United States (ranging from West Virginia up to the Canadian border on the east, and extending as far west as Minnesota), southeastern areas of Ontario, southeast to western regions of Manitoba, as well as in the southwestern reaches of Québec (Figure 2). There have been 11 reports of this species in Saskatchewan, with one confirmed and two probable cases of breeding from the far southeast (Smith 1996); the species is currently considered "Accidental" in this province (J. Pepper, pers. comm.). However, given the ongoing range expansion it is likely that this number will increase in coming years.




Figure 2. Breeding range of the Golden-winged Warbler (Dunn and Garrett 1997).

Wintering Golden-winged Warblers are found in areas of Central America (Guatemala, Honduras, Nicaragua, Costa Rica and Panama; Figure 3). They appear to be found in quantity in northern South America (Colombia, Venezuela), and in southern

Central America, with smaller numbers in the Greater Antilles (Cuba) and on some of the Caribbean islands (Hilty 1980; Johnson 1980; Moore 1980; Orejuela *et al.* 1980; Raffaele 1989; Raffaele *et al.* 1998; Ridgely and Gwynne 1989).

Figure 3. Wintering range of the Golden-winged Warbler (Based on Rappole *et al*

Québec – After moving northward to New England and Ontario, Golden-winged Warblers became established in extreme southern Québec in the early 1970s. While never very abundant, numbers were thought to peak in the late-1980s to early-1990s (N = 19 territorial males in one location) after which time numbers started to decrease; in 2001 the species was only found in 9 locations in the province (Gauthier and Aubry 1996; Environment Canada – CWS Québec Region)

To date Golden-winged Warblers have been reported on five Breeding Bird Survey routes in southern Québec (Sauer *et al.* 2005) and have been recorded in 31 of 2,464 blocks (1.3%) in an atlas project completed in the 1990s (Gauthier and Aubry 1996). Québec currently supports 0.2% of the global population of Golden-winged Warblers (P. Blancher, pers. comm.; Sauer *et al.* 2005).

Saskatchewan – The Saskatchewan Conservation Data Centre report 11 cases of Golden-winged Warblers in Saskatchewan (two from Saskatoon, three from Regina and the remaining six from the east side of the province) to date, and as such it is considered an accidental bird in the province (J. Keith, pers. comm.; Saskatchewan Bird Atlas Project; Smith 1996; J. Pepper, pers. comm.). The first record was from 1962 and they have appeared sporadically since that time with the latest record in 1999 (Saskatchewan Bird Atlas Project). No Golden-winged Warblers have been reported on Breeding Bird Surveys.

One case of breeding has been confirmed in the province and there are several reports of probable breeding in the far southeast of the province (Smith 1996; Saskatchewan Bird Atlas Project). The most recent records come from Duck Mountain near the Saskatchewan-Manitoba border.

## HABITAT

### Habitat requirements

#### a) Breeding grounds

Golden-winged Warblers aggressively defend all purpose territories from neighbouring conspecifics. The average territory ranges in size from 1-2 hectares and is considered large relative to that of other warbler species. Golden-winged Warblers tend to nest in loose aggregations or “colonies” that typically contain up to ten pairs of breeding birds (Confer and Knapp 1981; R. Fraser, unpub. data). Generally, areas that are logged, burned, and intermittently farmed can readily support the species.

Territories tend to contain patches of herbs and low shrubs (used for nests which are placed on the ground) and scattered trees plus a forested edge used for song posts and foraging (Frech and Confer 1987). Examples of some preferred habitat areas include hydro/utility right-of-ways, field edges, recently logged areas and beaver marshes. On the Canadian Shield alder bogs are often used, especially when there are some tall tree species present including Black Ash (*Fraxinus nigra*; Mills 1987).

The Golden-winged Warbler is a habitat specialist given its reliance on areas of early successional scrub (10-30 years into succession); they will not persist in an area when the stage of succession has exceeded their requirements (Confer and Knapp 1981; Confer and Larkin 1998). Territories in the earliest stages of succession, with high herb cover, support larger clutches than areas with higher tree and shrub cover (Confer *et al.* 2003)

#### b) Migration

Very little is known about Golden-winged Warbler habitat use while on migration; they appear to use various open woodland habitats, pine-oak, and scrub, often in foothill regions (NatureServe 2004) likely due to their reliance on insectivorous food items such as Lepidopteran larvae.

#### c) Wintering grounds

Golden-winged Warblers have been observed in mixed species flocks of Neotropical passerines on the wintering grounds in Costa Rica (Buehler *et al.* 2003), in Panama (Moore 1980) and in Colombia (Johnson 1980). These flocks require areas of open woodland and can be found in evergreen and semi-deciduous forests (Stiles *et al.* 1991). Golden-winged Warblers are rarely found in mixed-species flocks with conspecifics or Blue-winged Warblers (with generally only one Golden-winged Warbler observed in such mixed-species flocks; Moore 1980; Tramer and Kemp 1980).

The species has been observed at high elevations (1,500-3,000 m) while on the wintering grounds in various types of open woodland habitats, pine-oak and scrub (Hilty *et al.* 1985; Raffaele *et al.* 1998; Ridgely and Gwynne, Jr. 1989; Stiles *et al.* 1991; Stotz *et al.* 1996). Exceptions to this were observed by Keast (1980) who defined Blue-winged and Golden-winged Warblers as lowland dwellers, but contrastingly, also found Golden-winged Warblers at 1,000-3,000 m in Venezuela, and also in the Chiriqui highland rainforest in Panama. Within the preferred habitat types, the species is found along edges, primarily in the canopy (Stiles *et al.* 1991).

### **Habitat trends**

Of the 29 avian species that have showed significant population declines between 1980 and 2000 (Breeding Bird Survey data - Sauer *et al.* 2005), 90% of these use disturbance-generated ecosystems. Such ecosystems can be broadly defined as consisting of open fields, shrublands, mid-successional forests, open parkland and forest edges (Confer and Pascoe 2003).

Before the arrival of Europeans scrubby habitat was created when native North Americans employed slash-and-burn agriculture (Askins 2000). After European arrival scrubland habitat likely declined due to intensive grazing and farming. However, beginning in the 1840s upwards of tens of millions of acres of farmland was abandoned once again creating areas of suitable habitat. Much of this land has now passed

through old-field-succession and has become reforested (Askins 1993, 2000; Confer and Pascoe 2003) leading to an overall decrease in the availability of this habitat type in eastern North America (Confer and Pascoe 2003; Dettmers 2003; Gill 2004). In addition, given extensive human development, restoration of historical habitat types is not possible in many areas. Active management of disturbance-generated habitats is crucial for the survival of numerous bird species in North America, including the Golden-winged Warbler (Askins 1993; Confer and Pascoe 2003).

### **Habitat protection/ownership**

Given their reliance on early successional scrub this species is often found on land that is privately owned. General estimates on the proportion of breeding Golden-winged Warblers found on public vs. private lands throughout Canada are unavailable.

Areas of protection in Canada include some privately owned land, parks owned by Parks Canada as well as by the Ontario Ministry of Natural Resources and the National Capital Commission (Gatineau). Data from Parks Canada show that Golden-winged Warblers have been seen in five national parks in Ontario: Bruce Peninsula/Fathom Five, Georgian Bay Islands, Point Pelee, St. Lawrence Islands and Pukaskwa (but does not breed at Point Pelee or Pukaskwa) as well as in Riding Mountain National Park in Manitoba (P. Achuff, pers. comm.). At least one university in Ontario (i.e. Queen's University; R.J. Robertson, pers. comm.) also protects and maintains large tracts of land used by this species.

## **BIOLOGY**

### **Life cycle and reproduction**

Breeding pairs raise only one brood per year with an average clutch size of 4.75 (range 2 – 6) in Ontario. Clutch sizes of subsequent broods (i.e. renests that occur after a failed nesting event) often decrease with two eggs being the minimum observed in a third nesting attempt within one breeding season. Double brooding has not been reported in this species. Birds often breed at one year of age and can continue to reproduce up to nine years of age. Average generation time for this species is two to three years (R. Fraser, unpub. data). Pairs are socially monogamous but exhibit high rates of extra-pair paternity (i.e. social fathers are not genetic fathers 33% of the time; Fraser *et al.*, under review)

In a sample of 103 nests found in Ontario between 2001 and 2004, 55.3% (N=57) of nests successfully fledged at least one offspring, 37.9% (N=39) were depredated, and 6.8% (N=7) were abandoned. Of 435 nestlings in these nests, 256 (58.85%) fledged, while 176 (40.46%) were depredated. Average fledging success was 2.49 nestlings per breeding pair in this population (R. Fraser, unpub. data), which is similar to success rates from New York (2.3 per breeding pair; Confer *et al.* 2003) but lower than reported fledgling success rates from other regions (e.g. 3.7 per breeding pair in



Tennessee; Klaus and Buehler 2001). Nests can be abandoned due to flooding, partial predation, Brown-headed Cowbird parasitism and death of the female (R. Fraser, unpub. data).

Birds are known to be strictly insectivorous during the breeding season. Diet consists mainly of tortricid moths and their larvae (Confer 1992). Other moths and their pupae, other winged insects, spiders and spider egg sacks are also sometimes consumed (R. Fraser, unpub. data). Similar feeding habits are also seen on the wintering grounds.

## **Predation**

some areas of the United States are much lower (e.g. 16% in West Virginia; R. Canterbury, pers. comm.) suggesting that patterns of high return rates are not to be interpreted as being a range wide phenomenon and are likely attributable a variety of factors such as changes to the breeding habitat and/or to the surrounding landscape.

## Interspecific interactions

### Hybridization with Blue-winged Warblers - Overview

One of the key factors implicated in Golden-winged Warbler decline is hybridization with the Blue-winged Warbler. While traditionally breeding in allopatry after glaciers receded, these sister species were brought into geographic contact in the 1800s when clearing of the land facilitated the movement of Blue-winged Warblers into the Golden-winged Warbler breeding range in the northeastern United States (Gill 1980). Therefore, hybridization between these two superspecies is a relatively recent phenomenon exacerbated by anthropogenic disturbances to the landscape (Mayr and Short 1970; Gill 1980). The current hybrid zone extends from the eastern reaches of the species' range through to Minnesota, and is rapidly moving northward (Figure 4).




Figure 4. Hybrid index indicating remaining areas of allopatry and areas of contact. Golden-winged Warbler in yellow, Blue-winged Warbler in blue, hybrid zone in green (Data courtesy of K. Rosenberg, GOWAP, Cornell Lab of Ornithology).

In areas of sympatry, hybridization occurs with great frequency and has been seen in all areas of contact (e.g. Gill 1980; Confer 1992). Hybridization tends to negatively affect the Golden-winged Warbler with local extirpation occurring within 50 years of Blue-winged Warbler arrival being the norm (Gill 1997), although replacement can occur within as few as four or five years (Gill 2004). Advancing Blue-winged Warbler females apparently lead the introgression causing an asymmetric and apparently rapid

introgression of Blue-winged Warbler DNA into Golden-winged Warbler populations (Gill 1997). These results suggest that Blue-winged Warblers have a genetic advantage over Golden-winged Warblers and this may explain the replacement of Golden-winged Warblers by Blue-winged Warblers in areas of contact.

Recent genetic analyses, however, indicate that there is bidirectional gene flow in at least five currently mixed populations (Michigan, Ohio, West Virginia, New York and Ontario) suggesting that Blue-winged Warblers do not genetically swamp Golden-winged Warblers in all areas of contact (Shapiro *et al.*, 2004; Dabrowski *et al.*, in press). These results are important because they suggest that patterns of mitochondrial introgression that vary by site may be related to differences in geographic variables, in local population sizes, as well as subtle between-species differences in mate-choice and habitat preferences.

Further work examining habitat use and genetic analyses are needed from sympatric and allopatric populations of these species throughout the entire breeding range to elucidate the true impact of hybridization on populations of Golden-winged Warblers.

#### Hybrid phenotypes

When Blue-winged and Golden-winged Warblers interbreed they tend to produce one of two phenotypes assigned the names “Brewster’s Warbler” (“*V. leucobronchialis*”) and “Lawrence’s Warbler” (“*V. lawrencei*”), although exceptions are sometimes observed. The “Brewster’s Warbler” exhibits plumage characteristic of a Golden-winged Warbler with a yellow wing patch and white underparts (although some have speculated that a first generation “Brewster’s Warbler” may exhibit yellow underparts; Parkes 1951), but they lack the striking black throat and eye patch (Figure 5). The “Lawrence’s Warbler” looks mostly like a Blue-winged Warbler with yellow underparts and white wing bars. However, the “Lawrence’s Warbler” exhibits the throat and eye patch characteristic of the Golden-winged Warbler (Figure 5), likely representing a combination of recessive characters (Parkes 1951).



Figure 5. Male “Lawrence’s Warbler” (left) and a male “Brewster’s Warbler” (right). (Photos by Rachel Fraser).

Later generations of hybrids may show only subtle signs of introgression and may be classified as one of the parental species without close (i.e. in-the-hand) examination (Short 1963, 1969). Furthermore, both hybrid phenotypes sing one of the two parental song types, and very rarely a combination of them both (Ficken and Ficken 1968; Gill and Murray 1972). The latter two factors could have pronounced implications for the use of surveys to identify/locate Golden-winged Warblers.

Some studies have suggested that hybrids are at a mating disadvantage and therefore may have overall lower fitness than do the parental species (Confer and Tupper 2000; Confer and Barker 2002). Recent quantitative analyses of reproductive success, however, show that hybrids are not at a disadvantage and in fact hybrid fitness and extra-pair fertilizations are likely playing a major role in the ongoing hybridization between Blue-winged and Golden-winged Warblers (Fraser *et al.* under review; Gill 2004).

### Genetic distinctiveness

Prior to the publication of results from genetic analyses it was suggested that Blue-winged and Golden-winged Warblers had a common ancestry that occurred during the recent glacial maxima at approximately 20,000 years before present (e.g., Short 1963). As such, and given the existence of extensive hybridization in many areas, some researchers suggested that Blue-winged and Golden-winged Warblers should be considered conspecific (Mayr and Short 1970).

However, Gill (1997) reported that the nucleotide divergence between these 2 species was estimated to be 3.2% based on RFLP analyses. This suggests these lineages are substantially older and more differentiated than was previously suggested (Short 1963; Shapiro *et al.* 2004). Later work by Dabrowski *et al.* (in press) and Shapiro *et al.* (2004) determined the level of mtDNA sequence divergence to be approximately 4.5%. This level of divergence between ancestral Blue-winged and ancestral Golden-winged Warbler haplotypes is several orders of magnitude greater than expected if these lineages split near the Pleistocene-Holocene boundary. In comparison with other passerine genera the level of nucleotide divergence between the ancestral Blue-winged and ancestral Golden-winged haplotype groups is equivalent to the separation of many other pairs of taxa that comprise clear biological species (reviewed in Johnson and Cicero 2004).

### Regions of allopatry

Currently allopatric populations of Golden-winged Warblers only occur in the most extreme northern reaches of the breeding range, as well as at its highest nesting elevations in the Appalachian Mountains (Confer and Knapp 1981; Figure 4). Genetic purity of these populations has not yet been established, but is currently under investigation (R. Fraser, unpub. data). Even populations that were thought to be “safe havens” are now showing evidence of hybrid or Blue-winged Warbler arrival. For example, a previously allopatric population of Golden-winged Warblers in Ontario saw

the first arrival of hybrids and Blue-winged Warblers in the late 1980s. By 2004 17% of the breeding population was made up of hybrid phenotypes (Blue-winged Warbler occurrence remains low at <1%; R. Fraser, unpub, data). In addition, there have already been five reports of Blue-winged Warblers and one hybrid from Saskatchewan (Saskatchewan Bird Atlas Project; J. Keith, pers. comm.), while Manitoba reports only one hybrid sighting to date (a female "Brewster's Warbler" in 1932; Manitoba Museum - MARC record #2963). One male Blue-winged Warbler was sighted in Manitoba BlueJ-0arch Chilittea

data represent the most comprehensive data set in North America for analysis of bird population trends.

The entire Golden-winged Warbler breeding range falls within the BBS coverage area. However, several caveats in using these data should be acknowledged: 1) in areas of hybridization between Blue-winged and Golden-winged Warblers, hybrids and phenotypic Blue-winged Warblers may also sing the Golden-winged Warbler song and visual confirmation is not required (Sauer *et al.* 2005); 2) later generation hybrids often show only subtle signs of introgression meaning that their classification would be, incorrectly, to the parental species; and 3) BBS routes are limited to areas with roads and experienced observers, leading to inadequate sampling of the species in some areas.

The Ontario Breeding Bird Atlas (OBBA) also conducts surveys to determine prevalence of Golden-winged Warblers in the province, using coverage of 10-km squares as well as random point counts. Data currently available from the second OBBA come from a search effort about 85% of that of the first atlas so comparisons made between the two should accurately reflect changes in abundance of the Golden-winged Warbler (M. Cadman, pers. comm.). However, the first OBBA (1981-1985) treated the Golden-winged/Blue-winged Warbler complex differently than the second (2001-2005). As with the BBS, “heard only” records were used for the first atlas, while now only records that actually involve a sighting of the singing bird are included. This makes comparisons between the two atlases more difficult (M. Cadman, pers. comm.) although the data are likely still indicative of population trends.

## **Abundance**

Estimates of population size are “best guesses” and are based primarily on BBS data which have limitations (discussed above) – actual numbers of Golden-winged Warblers in Canada are yet to be confirmed and this can only be accomplished by intensive surveys in a wide variety of terrains and locations.

The global population size is estimated at 105,000 to 270,000 breeding pairs (K. Rosenberg, pers. comm.; Partners in Flight (PIF) data; Sauer *et al.* 2005). Forty percent of the population is thought to breed in Minnesota, with 69% of all the birds being found in Minnesota, Michigan and Wisconsin. Canada currently supports an estimated 18.5% of the breeding population (N = 19,425 - 49, 950), with 18.2% of the global population being found in Ontario (N = 19,110 - 49,140; P. Blancher, pers. comm.; BBS data, Sauer *et al.* 2005).

## **Fluctuations and trends**

Declines have been taking place for over 30 years (Gill 1997; Sauer *et al.* 2005) with the current rate of decline being estimated at 3.4%/year (P = 0.00001, 241 routes) in the United States, with declines as steep as 8.5% /year (P = 0.000, 127 routes) being seen in USFWS Region 5 (see Table 2 for list of States; Sauer *et al.* 2005). The species

is currently experiencing an overall (i.e. U.S. and Canada combined) decline of 2.4%/year ( $P = 0.001$ , 271 routes), making the Golden-winged Warbler one of the most vulnerable and steeply declining of all North American passerines.

In Canada, BBS data between 1968 and 2002 indicate a significant annual increase of 8.9% ( $p < .05$ , 47 routes), but there has been a dramatic 79% decrease in the last ten years (-14.4%/year,  $P < .05$ , 37 routes, 1993-2002; CWS 2005).

An Ontario Breeding Bird Atlas is currently in the final year of data collection (i.e. 2005) which will provide an up-to-date picture of what the trends are for this species in Ontario. In the first Ontario BBA (1981-1985) Golden-winged Warblers were located in 469 10-km squares, whereas data collected for the second atlas thus far (2001-2004) reports them in 278 squares. Their range has thought to have contracted in 200 squares (i.e. were found in these squares in the first atlas, but were not located in the same squares during the second atlas), and has expanded in 118 squares. There is an overlap between the first and second atlases in 160 squares (34%).

In contrast, the Blue-winged Warbler occurred in 139 squares in the first atlas, and has been recorded in 196 squares in the second atlas thus far. In addition, while the Blue-winged Warbler has contracted its range in 47 squares, it has expanded in 123 squares. These data suggest that Golden-winged Warblers are decreasing in many areas of Ontario, while Blue-winged Warblers and their hybrids are increasing (BBA data supplied by M. Cadman, pers. comm.).

Banding data from the Long Point Bird Observatory (LPBO, Port Rowan, ON) show that the number of captured Golden-winged Warblers increased until the mid-1980s then decreased (Table 1). The number of Blue-winged Warblers banded at LPBO has also been increasing over the past thirty years, and at a much greater rate than Golden-winged Warblers. Indeed, there has been a change in the ratio of Golden-winged Warblers to Blue-winged Warblers from 2.4:1 to 0.14:1 over a period of thirty years (LPBO data; J. McCracken, pers. comm.; Table 1).

**Table 1. Banding data from Long Point Bird Observatory (LPBO) for Golden-winged (GWWA) and Blue-winged Warblers (BWWA) from 1960-2004 (LPBO data provided by J.D. McCracken).**

9-year period	Number Banded		Ratio (GWWA:BWWA)
	GWWA	BWWA	
1960-1968	12	5	2.4:1
1969-1977	12	8	1.5:1
1978-1986	35	65	0.54:1
1987-1995	61	182	0.34:1
1996-2004	25	173	0.14:1

In summary, Golden-winged Warbler numbers are declining overall. Declines are less pronounced in Canada than they are in the U.S., but the rate of decline is now increasing in many areas of the country, especially in Ontario. Some areas may still be experiencing increases associated with their range expansion (e.g. Saskatchewan and Manitoba) but this is not necessarily indicative of them having greater success in these areas than in others and is likely associated with, at least in part, their allopatric status.

### **Rescue effect**

The ongoing range expansion of Golden-winged Warblers into Canada means that populations will likely continue to increase in some areas. However, whether these populations can sustain the species in Canada will depend on both habitat availability and arrival of Blue-winged Warblers. The Golden-winged Warbler may persist in refugia at higher elevations and latitudes, and perhaps in swamps where contact with Blue-winged Warblers is limited (Gill 2004).

## **LIMITING FACTORS AND THREATS**

Golden-winged Warbler declines are likely due to three main pressures they face while on the breeding grounds: a decrease in early successional scrub environments required for breeding, hybridization with the Blue-winged Warbler and perhaps Brown-headed Cowbird parasitism.

Of these three threats, loss of breeding habitat and hybridization with Blue-winged Warblers are the most significant contributors to Golden-winged Warbler declines. However, further research is needed to elucidate the impacts of these threats in concert, or to determine which one is having a more pronounced impact on Golden-winged Warbler populations in Canada.

### **a) Habitat loss**

Due to their reliance on anthropogenic disturbances (e.g. logging, hydro right-of-ways, and abandoned farmland) the Golden-winged Warbler is vulnerable to local extirpation throughout the breeding range. Declines occur with advancing succession and reforestation (Confer 1992) and a general loss of early successional scrub habitat in eastern North America (Confer and Pascoe 2003; Dettmers 2003). As such, it is likely that habitat loss is playing a major role in Golden-winged Warbler declines throughout the breeding range. This is further supported from across-species data showing that the majority of avian species undergoing population declines in North America depend on disturbance-generated ecosystems (Confer and Pascoe 2003; Sauer *et al.* 2005).

However, habitat limitations cannot be solely responsible for Golden-winged Warbler declines as they sometimes are extirpated from areas where suitable habitat remains, but only when sympatric with Blue-winged Warblers, suggesting that interactions between these species are also playing a role in declines (J. Confer, pers. comm.)



## **b) Hybridization with Blue-winged Warblers**

As detailed above (see Interspecific Interactions), hybridization between Blue-winged and Golden-winged Warblers is ongoing and widespread, and replacement of Golden-winged Warblers after 50 years of Blue-winged Warbler arrival is the norm (Gill 1997). Only one region of sympatry has persisted for more than 100 years and reasons for this coexistence are unknown (J. Confer, pers. comm.; Confer 2000).

At present there is no clear genetic r  
ss

etpal.

-1.

Tm0(18

**Table 2. State summaries of legal status of Golden-winged Warbler; it is unlisted in states not shown. Adapted from U.S. Golden-winged Warbler draft status assessment (Buehler *et al.*, 2003).**

REGION	STATE	DESIGNATION
USFWS Region 3	Indiana	State Endangered
	Ohio	State Endangered
Tenn	Wisconsin	Special Concern
nt	Special(Con)	Special Concern
g	6(c34)	Special Concern
	55637	Rared
Vermont		

)JTJ-0.0002 USFWS(Re

<b>State/Province</b>	<b>State Rank (season)</b>
New Jersey	3 (breeding)
New Mexico	1 (non-breeding)
New York	4
North Carolina	3 (breeding)
North Dakota	3
Ohio	1
Oklahoma	NR (non-breeding)
Pennsylvania	4 (breeding)
Rhode Island	X (breeding)
	2 (non-breeding)
Tennessee	3 (breeding)
Texas	3
Vermont	2-3 (breeding)
Virginia	3 (breeding)
West Virginia	3 (breeding)
	3 (non-breeding)
Wisconsin	4

1=Critically imperiled because of extreme rarity (often 5 or fewer occurrences) or because of some factor(s) such as very steep declines making it especially vulnerable to extirpation from the state or province, 2=Imperiled because of rarity due to very restricted range, very few populations (often 20 or fewer), steep declines, or other factors making it very vulnerable to extirpation, 3=Vulnerable due to a restricted range, relatively few populations (often 80 or fewer), recent and widespread declines, or other factors making it vulnerable to extirpation, 4=Uncommon but not rare; some cause for long-term concern due to declines or other factors, 5=Common, widespread, and abundant, NR=unranked, X=extirpated; states where ranked as NA (Not applicable because not suitable for conservation activities) are not listed.

## TECHNICAL SUMMARY

***Vermivora chrysoptera***

Golden-winged Warbler

Paruline à ailes dorées

Range of Occurrence in Canada: Manitoba, Ontario, Québec

---

Extent and Area Information

-

### Status and Reasons for Designation

**Status:** Threatened

**Alpha-numeric code:** Met criterion for Endangered, A2be, but designated Threatened because the species is still widespread, shows the ability to maintain small pure populations within the Blue-winged Warbler range, is still expanding in Manitoba, and is thus not in imminent danger of extinction. Criterion met for

## **ACKNOWLEDGEMENTS AND AUTHORITIES CONTACTED**

Thanks to the official sponsors of the Ontario Breeding Bird Atlas (Bird Studies Canada, Canadian Wildlife Service, Federation of Ontario Naturalists, Ontario Field Ornithologists, and Ontario Ministry of Natural Resources) for supplying Atlas data, and to the thousands of volunteer participants who gathered data for the project.

Thanks to Bird Studies Canada for providing data from the Long Point Bird Observatory's migration monitoring program, and to the countless volunteers and staff who have gathered and computerized these data over the years. The migration monitoring program receives important support from the Ontario Ministry of Natural Resources' Wildlife Assessment Program.

Thanks to Raleigh J. Robertson, Dick Cannings, Mary Stapleton and Kevin Fraser for providing valuable input and guidance during the compilation of this report, as well as to all the individuals and organizations who provided information and data; without your help this would not have been possible.

Funding for this report was provided by the Canadian Wildlife Service, Environment Canada.

### **A. Federal Representatives**

Achuff, P.L. October 2004. National Botanist, Ecological Integrity Branch, Parks Canada, Waterton Lakes National Park, Waterton Park, Alberta, T0K 2M0

Blancher, P. October 2004. Partners in Flight Scientist, Bird Studies Canada, P.O. Box 160, 115 Front Street, Port Rowan, Ontario N0E 1M0

Goulet, G. October 2004. Aboriginal Traditional Knowledge Coordinator, COSEWIC Secretariat, Canadian Wildlife Service, Environment Canada, Ottawa, Ontario K1A 0H3

Lougheed, C. October 2004. Scientific Project Officer, COSEWIC Secretariat, Canadian Wildlife Service, Environment Canada, Ottawa, Ontario K1A 0H3

McKay, Vicki. November 2004. Species at Risk Biologist, Point Pelee National Park of Canada, Parks Canada Agency, 407 Monarch Lane, R.R. #1, Leamington, Ontario N8H 3V4

### **B. Provincial/Territorial Jurisdictions**

#### Manitoba

Manitoba Conservation Data Centre. September 2004. P.O. Box 24, 200 Saulteaux Crescent, Winnipeg, Manitoba R3J 3W3

Duncan, J. October 2004. Manager, Biodiversity Conservation Section, Wildlife and Ecosystem Protection Branch, Manitoba Conservation, Box 24, 200 Saulteaux Crescent, Winnipeg, Manitoba R3J 3W3

Klapecki, J. September 2004. Collection Manager, Natural History Division, The Manitoba Museum, 190 Rupert Avenue, Winnipeg, Manitoba R3B 0N2  
Koes, R. September 2004. Manitoba Avian Research Committee, c/o Manitoba Naturalists Society, 401-63 Albert Street, Winnipeg, Manitoba R3B 1G4  
Lindgren, C. August 2004. Provincial Coordinator, Ducks Unlimited, Oak Hammock Marsh, P.O. Box 1160, Stonewall, Manitoba R0C 2Z0  
Tarleton, P. October 2004. Manager, Resource Conservation, Riding Mountain National Park of Canada, Wasagaming, Manitoba.

## Ontario

Cadman, M. August 2004. Coordinator, Ontario Breeding Bird Atlas, c/o University of Guelph, 50 Stone Road East, Blackwood Hall Room 211, Guelph, Ontario N1G 2W1  
Dextrase, A. September 2004. Senior SAR Biologist, Species At Risk Section, Ontario Parks, Ontario Ministry of Natural Resources, P.O. Box 7000, Peterborough, Ontario K9J 8M5  
Fraser, K.C. August 2004. Master's Student, Department of Biology, Queen's University, Kingston, Ontario K7L 3N6  
MacKenzie, J. October 2004. Coordinator, Ontario Natural Heritage Information Centre, 300 Water Street, 2nd Floor, North Tower, Peterborough, Ontario K9J 8M5  
McCracken, J. August 2004. Ontario Program Manager, Bird Studies Canada, P.O. Box 160, 115 Front Street, Port Rowan, Ontario N0E 1M0  
Promaine, A. October 2004. Park Ecologist. Georgian Bay Islands National Park, Parks Canada, 901 Wye Valley Road, Midland, Ontario L4R 4K6  
Robertson, R.J. July 2004. Professor of Biology and Baillie Family Chair in Conservation Biology, Department of Biology, Queen's University, Kingston, Ontario K7L 3N6  
Sutherland, D. September 2004. Natural Heritage Zoologist, Ontario Natural Heritage Information Center, Ontario Ministry of Natural Resources, 300 Water Street, 2<sup>nd</sup> Floor N., P.O. Box 7000, Peterborough, Ontario K9J 8M5  
Wade, K. October 2004. Ecosystem Scientist II, Pukaskwa National Park, Parks Canada. P.O. Box 212, Heron Bay, Ontario P0T 1R0

## Québec

Centre de données sur le patrimoine naturel du Québec, Ministère de l'Environnement du Québec, Direction de la conservation et du patrimoine écologique, 675 boulevard René-Lévesque Est – 10<sup>e</sup> étage, Québec, Québec G1R 5V7  
Banville, Daniel. October 2004. Coordonnateur espèces menacées et vulnérables, Ministère des Ressources naturelles, de la Faune et des Parcs du Québec, Faune Québec, Direction du développement de la faune, Édifice Marie-Guyard, 11<sup>e</sup> étage, 675, boul. René-Lévesque Est, Québec, Québec G1R 5V7

## Saskatchewan

- Hobson, K. October 2004. Researcher. Canadian Wildlife Service, Environment Canada, 115 Perimeter Road, Saskatoon, Saskatchewan S7N 0X4
- Keith, J. October 2004. Information Manager, Saskatchewan Conservation Data Centre, Resources Stewardship Branch, Saskatchewan Environment, Government of Saskatchewan, 3211 Albert Street, Regina, Saskatchewan S4S 5W6
- Pepper, J. October 2004. Saskatchewan Conservation Data Centre, Resources Stewardship Branch, Saskatchewan Environment, Government of Saskatchewan, 3211 Albert Street – Room 436, Regina, Saskatchewan S4S 5W6
- Van Wilgenburg, S. October 2004. Canadian Wildlife Service, Environment Canada, 115 Perimeter Road, Saskatoon, Saskatchewan S7N 0X4.

## INFORMATION SOURCES

- Achuff, P. pers. comm. 2004. *Email correspondence*. October 2004. National Botanist, Ecological Integrity Branch, Parks Canada, Waterton Lakes National Park, Waterton Park, Alberta.
- Askins, R.A. 1993. Population trends in grassland, shrubland, and forest birds in eastern North America. *Current Ornithology* 11:1-34.
- Askins, R.A. 2000. *Restoring North America's Birds – Lessons from landscape ecology*. Yale University Press, New Haven. pp. 39.
- Barker, S. 2005 *Email correspondence*. June 2005. GOWAP coordinator, Cornell Lab of Ornithology, Ithaca, NY.
- Berger, R. 2004. *Email correspondence*. M.N.R.M., Principal. Wildlife Resource Consulting Services MB Inc., Winnipeg, MB.
- BirdLife International. 2004. [Web resource]  
[http://www.birdlife.org/news/news/2004/10/state\\_us\\_birds.html](http://www.birdlife.org/news/news/2004/10/state_us_birds.html)
- Blancher, P. pers. comm. 2004. *Email correspondence*. October 2004. Partners in Flight Scientist, Bird Studies Canada, Port Rowan, Ontario.
- Buehler, D.A., R.A. Canterbury, and J.L. Confer. 2003. Golden-winged Warbler (*Vermivora chrysoptera*) Draft Status Assessment. Technical report prepared for the US Fish and Wildlife Service.
- Cadman, M. pers comm. 2004. *Email correspondence*. September 2004. Coordinator, Ontario Breeding Bird Atlas, Guelph, Ontario.
- Canterbury, R.A. pers. comm. 2000-2004. *Email correspondence*. October 2004. Professor of Biology, Concord University, Athens, West Virginia.
- Confer, J.L. pers. comm. 2000-2004. *Email correspondence and discussion*. August 2004. Professor of Biology, Ithaca College, Ithaca, New York.
- Confer, J.L. 1992. Golden-winged Warbler (*Vermivora chrysoptera*). *In* The Birds of North America, No. 20 (A. Poole, P. Stettenheim, and F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA.
- Confer, J.L. 2000. A reassessment of the status of Golden-winged and Blue-winged Warblers in the Hudson Highlands of Southern New York. *Wilson Bulletin* 112(4):544-546.



- Confer, J.L. and S.E. Barker. 2002. A tale of two species inscribed in DNA. *Birdscope: Newsletter of the Cornell Lab of Ornithology* 16(1). Available at:  
<http://www.birds.cornell.edu>
- Confer, J.L. and K. Knapp. 1981. Golden-winged Warblers and Blue-winged Warblers: The relative success of a habitat specialist and a generalist. *Auk* 98:108-114.
- Confer, J.L. and J.L. Larkin. 1998. Behavioural interactions between Golden-winged and Blue-winged Warblers. *Auk* 115(1):209-214.
- Confer, J.L., J.L. Larkin, and P.E. Allen. 2003. Effects of vegetation, interspecific

- Fraser, R., V.L. Friesen and R.J. Robertson. Under review. Paternity and performance of Golden-winged Warblers (*Vermivora chrysoptera*) and Golden-winged X Blue-winged Warbler (*V. pinus*) hybrids: implications of heterospecific mate choice. Submitted to Behavioral Ecology and Sociobiology.
- Frech, M. and J.L. Confer. 1987. The Golden-winged Warbler: Competition with the Blue-winged Warbler and habitat selection in portions of southern, central and northern New York. *The Kingbird* 37:65-71.
- Gauthier, J. and Y. Aubry (Eds). 1996. The breeding birds of Québec. Association québécoise des groupes d'ornithologues. The Province of Québec Society for the Protection of Birds and the Canadian Wildlife Service, Environment Canada. Québec Region.
- Gill, F.B. pers. comm. 2000-2004. *Email correspondence and conversation*. 2000-present. Chief Science Officer, National Audubon Society, Ivyland, PA.
- Gill, F.B. 1980. Historical aspects of hybridization between Blue-winged and Golden-winged Warblers. *Auk* 97(1):1-18.
- Gill, F.B. 1997. Local cytonuclear extinction of the Golden-winged Warbler. *Evolution* 51(2):519-525.
- Gill, F.B. 2004. Blue-winged Warblers (*Vermivora pinus*) versus Golden-winged Warblers (*V. chrysoptera*). *Auk* 121:1014-1018.
- Gill, F.B. and B.G. Murray, Jr. 1972. Discrimination behaviour and hybridization of the Blue-winged and Golden-winged Warblers. *Evolution* 26:282-293.
- Golden-winged Warbler Atlas Project (GOWAP): a citizen science project of the Cornell Laboratory of Ornithology <http://birds.cornell.edu/gowap/birds.html>
- Goulet, G. pers. comm. 2004. *Email correspondence*. October 2004. Aboriginal Traditional Knowledge Coordinator, COSEWIC Secretariat, Canadian Wildlife Service, Environment Canada, Ottawa, Ontario.
- Hilty, S.L. 1980. Relative abundance of north Temperate Zone breeding migrants in western Colombia and their impact on fruiting trees. Pp. 265-271 in Keast, A., E.S. Morton. 1980. *Migrant Birds in the Neotropics: Ecology, behavior, distribution, and conservation*. Smithsonian Inst. Press, Washington, DC.
- Hilty, S.L., W.L. Brown, and G. Tudor. 1985. *A Guide to the Birds of Colombia*. Princeton University Press, Princeton, NJ.
- Johnson, T.B. 1980. Resident and North American migrant bird interactions in the Santa Marta Highlands, northern Colombia. Pp. 239-247 in Keast, A., E.S. Morton. 1980. *Migrant Birds in the Neotropics: Ecology, behavior, distribution, and conservation*. Smithsonian Inst. Press, Washington, DC.
- Johnson, N.K. and C. Cicero. 2004. New mitochondrial DNA data affirm the importance of Pleistocene speciation in North American birds. *Evolution* 58:1122-1130.
- Keast, A. 1980. Spatial relationships between migratory parulid warblers and their ecological counterparts in the Neotropics, Pp. 109-130 in *Migrant birds in the Neotropics/ecology, behaviour, distribution and conservation* (A. Keast and E.S. Morton, Eds). Smithsonian Inst. Press, Washington, DC.
- Keith, J. pers. comm. 2004. *Email correspondence*. October 2004. Information Manager, Saskatchewan Conservation Data Centre, Resources Stewardship Branch, Saskatchewan Environment, Government of Saskatchewan, Regina, Saskatchewan.

- Klapecki, J. pers. comm. 2004. *Email correspondence*. September 2004. Collections Manager, Natural History Division, The Manitoba Museum, Winnipeg, Manitoba.
- Klaus, N.A. and D.A. Buehler. 2001. Golden-winged Warbler breeding habitat characteristics and nest success in clearcuts in the southern Appalachian Mountains. *Wilson Bulletin* 113:291-301.
- Klein, N.K., K.J. Burns, S.J. Hackett and C.S. Griffiths. 2004. Molecular phylogenetic relationships among the Wood Warblers (Parulidae) and historical biogeography in the Caribbean Basin. *Journal of Caribbean Ornithology* 17:3-17.
- Long Point Bird Observatory, Bird Studies Canada. 2004. Data provided by J.D. McCracken via email.
- Lovette, I.L. pers. comm. 2002-2004. *Discussion*. 2002-present. Director, Evolutionary Biology Program, Cornell Laboratory of Ornithology, Ithaca, NY.
- L.P. Canada Ltd. Swanvalley and Rob Berger. 2004. Supplied Manitoba survey data from Duck Mountain, Manitoba.
- McCracken, J.D. 2004. Ontario Program Manager, Bird Studies Canada. Port Rowan, ON.
- Manitoba Avian Research Committee. 2003. *The Birds of Manitoba*. Manitoba Naturalists Society, Winnipeg, Manitoba. VIII. 504 pp.
- Manitoba Museum. 2004. *Manitoba Bird Record Cards (1905-1992)*.
- Mayr, E. and L.L. Short. 1970. Species taxa of North American birds: A contribution of comparative systematics. *Nuttall Ornithological Club*, No. 9, 127 pp.
- Mills, A. 1987. Golden-winged Warbler. Pp. 358-359 in Cadman, M.D., P.E.J. Eagles, and F.M. Helleiner, eds. *Atlas of the breeding birds of Ontario*. University of Waterloo Press, Waterloo, ON.
- Moore, E.S. 1980. Adaptations to Seasonal Changes by Migrant Land Birds in the Panama Canal Zone. Pp 437-456 in Keast, A., E.S. Morton. 1980. *Migrant Birds in the Neotropics: Ecology, behavior, distribution, and conservation*. Smithsonian Inst. Press, Washington, D. C.
- NatureServe. 2004. NatureServe Explorer: An online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>.
- Ontario Breeding Bird Atlas. 2004. [Web application] <http://www.birdsontario.org/atlas/atlasmain.html>
- Orejuela, J.E., R.J. Raitt, and H. Alvarez. 1980. Differential use by North American migrants of three types of Colombian forests. Pp. 253-264 in Keast, A. and E.S. Morton, eds. *Migrant birds in the Neotropics: Ecology, behavior, distribution, and conservation*. Smithsonian Institution Press, Washington, D.C.
- Parkes, K.C. 1951. The genetics of the Golden-winged x Blue-winged Warbler complex. *Wilson Bulletin* 63(1):4-15.
- Partners in Flight (PIF), *Watch list for threatened and declining birds*, [Web application] [http://www.abcbirds.org/pif/pif\\_watch\\_list.htm](http://www.abcbirds.org/pif/pif_watch_list.htm)
- Partners in Flight (PIF) [Web application] <http://www.partnersinflight.org/>
- Pepper, J. 2004. *Email correspondence*. Resource Stewardship, Saskatchewan Environment, Regina, SK.
- Pyle, P. 1997. *Identification Guide to North American Birds*. Slate Creek Press, Bolinas, California. 732 pp.

- Raffaele, H. 1989. *A guide to the birds of Puerto Rico and the Virgin Islands*. Princeton University Press, Princeton, NJ.
- Raffaele, H., J. Wiley, O. Garrido, A. Keith, and J. Raffaele. 1998. *A guide to the birds of the West Indies*. Princeton University Press, Princeton, NJ.
- Rappole, J.H., E.S. Morton, T.E. Lovejoy, and J.L. Ruos. 1983. *Nearctic avian migrants in the Neotropics*. U.S. Fish and Wildlife Service.
- Ridgely, R.S. and J.A. Gwynne, Jr. 1989. *A guide to the birds of Panama, revised*. Princeton University Press, Princeton, NJ.
- Ridgely, R.S. and G. Tudor. 1989. *The birds of South America, Vol. 1*. University of Texas Press, Austin, Texas.
- Robertson, R.J. pers. comm. 2000-2005. *Email correspondence*. October 2004. Professor of Biology, and Baillie Family Chair in Conservation Biology, Queen's University, Kingston, Ontario.
- Rosenberg, K.V. pers. comm. 2002-2004. *Email correspondence and discussion*. August 2004. Partners in Flight Northeast Regional Coordinator, Director – Conservation Science Program, Cornell Laboratory of Ornithology, Ithaca, New York.
- Saskatchewan Bird Atlas Project. 2004. [Web application]  
<http://gisweb1.serm.gov.sk.ca/BirdAtlas/mainscreen.htm>
- Sauer, J.R., J.E. Hines, and J. Fallon. 2005. *The North American Breeding Bird Survey, Results and Analysis 1966 - 2004. Version 2005.2* [Web application]  
<http://www.pwrc.usgs.gov/>. Laurel, MD
- Shapiro, L.H., R.A. Canterbury, D.M. Strover, and R.C. Fleisher (2004) Reciprocal introgression between Golden-winged Warblers (*Vermivora chrysoptera*) and Blue-winged Warblers (*V. pinus*) in eastern North America. *Auk* 121:1019-1030.
- Short, Jr. L.L. 1963. Hybridization in wood warblers *Vermivora pinus* and *V. chrysoptera*. *Proceedings of the XIII International Ornithological Congress*:147-160.
- Short, Jr. L.L. 1969. "Isolating mechanisms" in the Blue-winged Warbler - Golden-winged Warbler complex. *Evolution* 23:355-356.
- Smith, A.R. 1996. *Atlas of Saskatchewan Birds*. Pp. 301. Saskatchewan Natural History Society, Nature Saskatchewan, Environment Canada.
- Social Science Research, Ontario Service Centre. Point Pelee National Park, Provincial Economic Impact Estimate for 2003-2004.
- Stiles, G.F., A.F. Skutch and D. Gardner. 1991. *A guide to the birds of Costa Rica*. Cornell Univ. Press, Ithaca NY. 511 pp.
- Stotz, D.F., J.W. Fitzpatrick, T.A. Parker, and D.K. Moskovits. 1996. *Neotropical Birds, Ecology and Conservation*. University of Chicago Press. 502 pp.
- Short, L.L. 1963. Hybridization in the Wood Warblers *Vermivora pinus* and *V. chrysoptera*. *Proceedings of the VIII International Ornithological Congress*: 147-160.
- Tarleton, P. pers. comm. 2004. *Email correspondence*. October 2004. Manager, Resource Conservation, Riding Mountain National Park of Canada, Wasagaming, Manitoba
- Tramer, E.J. and T.R. Kemp. 1980. Foraging ecology of migrant and resident warblers and vireos in the highlands of Costa Rica, Pp. 285-296 in *Migrant birds in the*

Neotropics: Ecology, behaviour, distribution, and conservation. (A. Keast and E.S. Morton, eds.). Smithsonian Inst. Press, Washington, DC.

Van Wilgenburg, S. pers. comm. 2004. *Email correspondence and provision of manuscript by Cumming, 1998*. October 2004. Canadian Wildlife Service, Environment Canada, 115 Perimeter Road, Saskatoon, Saskatchewan S7N 0X4.